

System Analytics

www.sa-journal.org

Syst. Anal. Vol. 2, No. 1 (2024) 35–47.

Paper Type: Original Article

On Min-Max Goal Programming Approach for Solving Piecewise Quadratic Fuzzy Multi-Objective De Novo Programming Problems

Hamiden Abd El- Wahed Khalifa^{1,*}, Seyed Ahmad Edalatpanah², Darko Bozanic³

¹ Department of Operations Research, Graduate School for Statistical Research, Cairo University, Giza, Egypt; hamiden@cu.edu.eg; Ha.Ahmed@qu.edu.sa.

² Department of Applied Mathematics, Ayendang Institute of Higher Education, Tonkabon, Iran; saedalat@yahoo.com & s.a.edalatpanah@aihe.ac.ir.

³ Associate professor, University of Defence in Belgrade, Military Academy, Belgrade, Serbia; darko.bozanic@va.mod.gov.rs; ddbozanic@gmail.com.

Citation:

Received: 14 February 2023

Revised: 25 April 2023

Accepted: 12 May 2023

Abd El- Wahed Khalifa, H., Edalatpanah, S. A., & Bozanic, D. (2024).

On min-max goal programming approach for solving piecewise quadratic fuzzy multi- objective de novo programming problems. *System Analytics*, 2(1), 35-47.

Abstract

De novo programming is considered an essential tool for establishing optimal system design. This paper studies the Multi-Objective De Novo Programming (MODNP) problem with Piecewise Quadratic Fuzzy (PQF) data in the objective function coefficients. One of the best interval approximations, namely, the close interval approximation of the PQF number, is applied to solve the MODNP problem. A necessary and sufficient condition for the solution from the efficiency standpoint is established. A Min-max goal programming approach with positive and negative ideals is proposed to obtain optimal compromise system design. The stability set of the first kind corresponding to the optimal system design is defined and determined. The stability set of the first kind corresponding to the optimal system design is determined. The steps of the proposed solution approach are illustrated through numerical examples.

Keywords: Multi-objective de novo programming, Piecewise quadratic fuzzy numbers, Close interval approximation, α -Fuzzy efficient, Goal programming, Optimal system design, Parametric study.

1 | Introduction

Multi-objective decision-making solution approaches have an essential role in decision problem solutions. According to the Decision Maker (DM) influence in the optimization process, Multi-Objective Optimization (MO) methods can be classified as [1]:

Corresponding Author: hamiden@cu.edu.eg

<https://doi.org/10.31181/sa21202411>



Licensee System Analytics. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0>).

- I. Methods where DM does not provide information (no-preference methods).
- II. Methods where a posteriori information is used (posterior methods).
- III. Methods where a priori information is used (priori methods).
- IV. Methods where progressive information is used (interactive methods).

Wang and Chaing [2] applied the user preference-enabling method to solve general constrained nonlinear MO problems. Kundu and Islam [3] introduced an interactive method to design a highly reliable and productive system with minimum cost to solve multi-objective reliability optimization problems. Waliv et al. [4] studied the effect of capital investment and warehouse space on profits and shortage cost through sensitivity analysis and compared the efficiency of fuzzy nonlinear programming and intuitionistic fuzzy optimization techniques to obtain the solution. Ahmed [5] proposed a method to solve MO problems with intuitionistic fuzzy parameters. Liu et al. [6] introduced a new systematic method for determining an optimal operation scheme for minimizing octane number loss and operational risks.

De Novo Programming (DNP) deals with the design of an optimal system design. Many researchers have studied the DNP problem (for instance, [7]–[13]).

Fuzzy sets theory, introduced by Zadeh [14], makes this possible. Fuzzy numerical data can be represented by employing fuzzy subsets of the real line, known as fuzzy numbers. Dubois [15] extended the use of algebraic operations on real numbers to fuzzy numbers by use of a fuzzification principle. Despite vast decision-making experience, the decision-maker cannot consistently articulate the goals precisely. Decision-making in a fuzzy environment, developed by Bellman and Zadeh [16], improved and was a great help in managing decision problems. Zimmermann [17] proposed the fuzzy set theory and its applications. Many approaches have been introduced for dealing with DNP problems [18], [19].

Goal Programming (GP) is one of the critical approaches in the multi-objective decision-making process, which is the extension of linear programming with the achievement of target objective values. Charnes and Cooper [20] first used the GP. Many authors applied GP in their research [20]–[23].

In his earlier work, Osman [25] introduced the notions of the solvability set, the stability set of the first kind, and the stability set of the second kind and analyzed these concepts for parametric convex nonlinear programming problems. Osman and El-Banna [26] studied the stability of multi-objective nonlinear programming problems with fuzzy parameters.

This paper introduces a Piecewise Quadratic Fuzzy Multi-Objective De Novo Programming (PQF-MODNP) problem with PQF data in the objective function coefficients. A GP approach is applied to obtain the optimal system design.

The remainder of the paper is organized as follows: Section 2 introduces some preliminaries needed in this paper. Section 3 formulates the mathematical model for a PQF MODNP problem. Section 4 characterizes the α – efficient solutions for *Problem (2)*. Section 5 investigates the goal-programming approach for obtaining optimal system design. Section 5 gives a numerical example for illustration. Finally, some concluding remarks are reported in Section 6.

2 | Preliminaries

To discuss the problem quickly, it recalls basic rules and findings related to fuzzy numbers, PQF numbers, close interval approximation, and its arithmetic operations.

Definition 1 ([14]). Fuzzy number: A fuzzy number $\tilde{A}$ is a fuzzy set with a membership function defined as $\pi_{\tilde{A}}(x): \mathfrak{R} \rightarrow [0,1]$, and satisfies:

- I. $\tilde{A}$ is fuzzy convex, i.e., $\pi_{\tilde{A}}(\delta x + (1 - \delta) y) \geq \min\{\pi_{\tilde{A}}(x), \pi_{\tilde{A}}(y)\}$; for all $x, y \in \mathfrak{R}$; $0 \leq \delta \leq 1$.
- II. $\tilde{A}$ is normal, i.e., $\exists x_0 \in \mathfrak{R}$ for which $\pi_{\tilde{A}}(x_0) = 1$.

III. $\text{Supp}(\tilde{A}) = \{x \in \mathbb{R}: \pi_{\tilde{A}}(x) > 0\}$ is the support of $\tilde{A}$.

IV. $\pi_{\tilde{A}}(x)$ is an upper semi-continuous (i. e., for each $\alpha \in (0,1)$, the α – cut set $\tilde{A}_\alpha = \{x \in \mathbb{R}: \pi_{\tilde{A}} \geq \alpha\}$ is closed.

Definition 2 ([27]). A Piecewise Quadratic Fuzzy Number (PQFN) is denoted by $\tilde{A}_{PQ} = (a_1, a_2, a_3, a_4, a_5)$, where $a_1 \leq a_2 \leq a_3 \leq a_4 \leq a_5$ are real numbers and are defined by whether their membership function $\mu_{\tilde{A}_{PQ}}$ is given by (see, Fig. 1).

$$\mu_{\tilde{A}_{PQ}} = \begin{cases} 0, & x < a_1, \\ \frac{1}{2} \frac{1}{(a_2 - a_1)^2} (x - a_1)^2, & a_1 \leq x \leq a_2, \\ \frac{1}{2} \frac{1}{(a_3 - a_2)^2} (x - a_3)^2 + 1, & a_2 \leq x \leq a_3, \\ \frac{1}{2} \frac{1}{(a_4 - a_3)^2} (x - a_3)^2 + 1, & a_3 \leq x \leq a_4, \\ \frac{1}{2} \frac{1}{(a_5 - a_4)^2} (x - a_5)^2, & a_4 \leq x \leq a_5, \\ 0, & x > a_5. \end{cases}$$




Fig. 1. Graphical Representation of a PQFN.

Definition 3 ([27]). Let $\tilde{A}_{PQ} = (a_1, a_2, a_3, a_4, a_5)$ and $\tilde{B}_{PQ} = (b_1, b_2, b_3, b_4, b_5)$ be two PQFNs. The arithmetic operations on $\tilde{A}_{PQ}$ and $\tilde{B}_{PQ}$ are.

I. Addition: $\tilde{A}_{PQ}(+) \tilde{B}_{PQ} = (a_1 + b_1, a_2 + b_2, a_3 + b_3, a_4 + b_4, a_5 + b_5)$.

II. Subtraction: $\tilde{A}_{PQ}(-) \tilde{B}_{PQ} = (a_1 + b_5, a_2 + b_4, a_3 + b_3, a_4 + b_2, a_5 + b_1)$.

III. Scalar multiplication: $k\tilde{A}_{PQ} = \begin{cases} (k a_1, k a_2, k a_3, k a_4, k a_5), & k > 0, \\ (k a_5, k a_4, k a_3, k a_2, k a_1), & k < 0. \end{cases}$

Definition 4 ([27]). An interval approximation $[A] = [a_{\alpha}^-, a_{\alpha}^+]$ of a PQFN $\tilde{A}$ is called closed interval approximation if

$a_{\alpha}^- = \inf\{x \in \mathbb{R}: \mu_{\tilde{A}} \geq 0.5\}$, and $a_{\alpha}^+ = \sup\{x \in \mathbb{R}: \mu_{\tilde{A}} \geq 0.5\}$.

Definition 5 ([27]). Let $[A] = [a_{\alpha}^-, a_{\alpha}^+]$, and $[B] = [b_{\alpha}^-, b_{\alpha}^+]$ be two interval approximations of PQFN. Then the arithmetic operations are.

I. Addition: $[A](+)[B] = [a_{\alpha}^- + b_{\alpha}^-, a_{\alpha}^+ + b_{\alpha}^+]$,

II. Subtraction: $[A](-)[B] = [a_{\alpha}^{-} - b_{\alpha}^{+}, a_{\alpha}^{+} - b_{\alpha}^{-}]$,

III. Scalar multiplication: $\alpha [A] = \begin{cases} [\alpha a_{\alpha}^{-}, \alpha a_{\alpha}^{+}], & \alpha > 0, \\ [\alpha a_{\alpha}^{+}, \alpha a_{\alpha}^{-}], & \alpha < 0. \end{cases}$

IV. Multiplication: $[A](\times)[B]$

$$\left[\frac{a_{\alpha}^{+} b_{\alpha}^{-} + a_{\alpha}^{-} b_{\alpha}^{+}}{2}, \frac{a_{\alpha}^{-} b_{\alpha}^{-} + a_{\alpha}^{+} b_{\alpha}^{+}}{2} \right],$$

V. Division: $[A](\div)[B]$

$$\begin{cases} \left[2 \left(\frac{a_{\alpha}^{-}}{b_{\alpha}^{-} + b_{\alpha}^{+}} \right), 2 \left(\frac{a_{\alpha}^{+}}{b_{\alpha}^{-} + b_{\alpha}^{+}} \right) \right], & [B] > 0, b_{\alpha}^{-} + b_{\alpha}^{+} \neq 0, \\ \left[2 \left(\frac{a_{\alpha}^{+}}{b_{\alpha}^{-} + b_{\alpha}^{+}} \right), 2 \left(\frac{a_{\alpha}^{-}}{b_{\alpha}^{-} + b_{\alpha}^{+}} \right) \right], & [B] < 0, b_{\alpha}^{-} + b_{\alpha}^{+} \neq 0. \end{cases}$$

VI. The order relations:

I. $[A](\lesssim)[B]$ if $a_{\alpha}^{-} \leq b_{\alpha}^{-}$ and $a_{\alpha}^{+} \leq b_{\alpha}^{+}$ or $a_{\alpha}^{-} + a_{\alpha}^{+} \leq b_{\alpha}^{-} + b_{\alpha}^{+}$.

II. $[A]$ is preferred to $[B]$ if and only if $a_{\alpha}^{-} \geq b_{\alpha}^{-}$, $a_{\alpha}^{+} \geq b_{\alpha}^{+}$.

It is noted that $P(\mathbb{R}) \subset F(\mathbb{R})$, where $F(\mathbb{R})$, and $P(\mathbb{R})$ are the sets of all PQFNs and close in interval approximation of PQFN, respectively.

3 | Problem Formulation

Consider a MODNP problem with PQF objective functions coefficients as

$$\begin{aligned} \max \tilde{Z}^p(x, \tilde{c}_{pj}) &= \sum_{j=1}^n \tilde{c}_{pj} x_j, p = \overline{1, P}, \\ \min \tilde{W}^q(x, \tilde{c}_{qj}) &= \sum_{j=1}^n \tilde{c}_{qj} x_j, q = \overline{1, Q}. \end{aligned}$$

Subject to

$$x \in X = \left\{ x^n \in \mathfrak{R}: \sum_{j=1}^n a_{ij} x_j - b_i \leq 0; \sum_{i=1}^m r_i x_i = B; x_j \geq 0, j = \overline{1, n} \right\}.$$

(1)

Where, $\tilde{c}_{pj}(p = \overline{1, P})$ and $\tilde{c}_{qj}(q = \overline{1, Q})$ are PQF variables on $\mathfrak{R}$ which are characterized by PQFNs., $x_j(j = \overline{1, n})$ and x_i are decision variables for projects and resources, respectively. r_i , represents the price of resource i , and B is the total availability budget.

It is noted that *Problem (1)* can be formulated as a continuous "knapsack" problem using the unit price of resource constraints

$$\begin{aligned} \max \tilde{Z}^p(x, \tilde{c}_{pj}) &= \sum_{j=1}^n \tilde{c}_{pj} x_j, p = \overline{1, P}, \\ \min \tilde{W}^q(x, \tilde{c}_{qj}) &= \sum_{j=1}^n \tilde{c}_{qj} x_j, q = \overline{1, Q}. \end{aligned}$$

Subject to

$$x \in X = \{x \in \mathfrak{R}^n: Vx \leq B; x_j \geq 0, j = \overline{1, n}\}.$$

(2)

Where, $Z^p = (Z^1, \dots, Z^P) \in \mathfrak{R}^P$, $W^q = (W^1, \dots, W^Q) \in \mathfrak{R}^Q$, $V = (V_1, \dots, V_n) = pA \in \mathfrak{R}^n$, B is the total given budget.

Definition 6 ([28]). $x^* \in X$ is an α – efficient solution for *Problem (2)* if there is no $x \in X$ such that:

$$\mu \left(\begin{array}{l} Z^1(x, \tilde{c}_{1j}) \geq Z^1(x^*, \tilde{c}_{1j}), \dots, Z^{p-1}(x, \tilde{c}_{p-1,j}) \geq Z^{p-1}(x^*, \tilde{c}_{p-1,j}), \\ Z^p(x, \tilde{c}_{pj}) \geq Z^p(x^*, \tilde{c}_{pj}), Z^{p+1}(x, \tilde{c}_{p+1,j}) \geq Z^{p+1}(x^*, \tilde{c}_{p+1,j}), \dots, \\ Z^p(x, \tilde{c}_{pj}) \geq Z^p(x^*, \tilde{c}_{pj}); W^1(x, \tilde{c}_{1j}) \leq W^1(x^*, \tilde{c}_{1j}), \dots, \\ W^{q-1}(x, \tilde{c}_{q-1,j}) \leq W^{q-1}(x^*, \tilde{c}_{q-1,j}), W^q(x, \tilde{c}_{qj}) \leq W^q(x^*, \tilde{c}_{qj}), \\ W^{q+1}(x, \tilde{c}_{q+1,j}) \leq W^{q+1}(x^*, \tilde{c}_{q+1,j}), \dots, W^Q(x, \tilde{c}_{Qj}) \leq W^Q(x^*, \tilde{c}_{Qj}) \end{array} \right) \geq \alpha. \quad (3)$$

On the account of the extension principle,

$$\mu \left(\begin{array}{l} Z^1(x, \tilde{c}_{1j}) \geq Z^1(x^*, \tilde{c}_{1j}), \dots, Z^{p-1}(x, \tilde{c}_{p-1,j}) \geq Z^{p-1}(x^*, \tilde{c}_{p-1,j}), \\ Z^p(x, \tilde{c}_{pj}) \geq Z^p(x^*, \tilde{c}_{pj}), Z^{p+1}(x, \tilde{c}_{p+1,j}) \geq Z^{p+1}(x^*, \tilde{c}_{p+1,j}), \dots, \\ Z^p(x, \tilde{c}_{pj}) \geq Z^p(x^*, \tilde{c}_{pj}); W^1(x, \tilde{c}_{1j}) \leq W^1(x^*, \tilde{c}_{1j}), \dots, \\ W^{q-1}(x, \tilde{c}_{q-1,j}) \leq W^{q-1}(x^*, \tilde{c}_{q-1,j}), W^q(x, \tilde{c}_{qj}) \leq W^q(x^*, \tilde{c}_{qj}), \\ W^{q+1}(x, \tilde{c}_{q+1,j}) \leq W^{q+1}(x^*, \tilde{c}_{q+1,j}), \dots, W^Q(x, \tilde{c}_{Qj}) \leq W^Q(x^*, \tilde{c}_{Qj}), \end{array} \right) = \quad (4)$$

$$\sup_{(c_1, \dots, c_p; c_{1j}, \dots, c_{Qj}) \in C \times D} \min \left(\begin{array}{l} \mu_{\tilde{c}_1}(c_1), \dots, \mu_{\tilde{c}_{p-1}}(c_{p-1}), \mu_{\tilde{c}_p}(c_p), \mu_{\tilde{c}_{p+1}}(c_{p+1}), \dots, \mu_{\tilde{c}_p}(c_p); \\ \mu_{\tilde{c}_1}(c_1), \dots, \mu_{\tilde{c}_{q-1}}(c_{q-1}), \mu_{\tilde{c}_q}(c_q), \mu_{\tilde{c}_{q+1}}(c_{q+1}), \dots, \mu_{\tilde{c}_Q}(c_Q) \end{array} \right).$$

Where

$$C = \left\{ \begin{array}{l} (c_1, \dots, c_p) \in \mathfrak{R}^{P(n)}: Z^1(x, c_{1j}) \geq Z^1(x^*, c_{1j}), \dots, \\ Z^{p-1}(x, c_{p-1,j}) \geq Z^{p-1}(x^*, c_{p-1,j}), Z^p(x, c_{pj}) \geq Z^p(x^*, c_{pj}), \\ Z^{p+1}(x, c_{p+1,j}) \geq Z^{p+1}(x^*, c_{p+1,j}), \dots, Z^p(x, c_{pj}) \geq Z^p(x^*, c_{pj}) \end{array} \right\}, \quad (5)$$

$$D = \left\{ \begin{array}{l} (c_1, \dots, c_Q) \in \mathfrak{R}^{P(n)}: W^1(x, c_{1j}) \leq W^1(x^*, c_{1j}), \dots, \\ W^{q-1}(x, c_{q-1,j}) \leq W^{q-1}(x^*, c_{q-1,j}), W^q(x, c_{qj}) \leq W^q(x^*, c_{qj}), \\ W^{q+1}(x, c_{q+1,j}) \leq W^{q+1}(x^*, c_{q+1,j}), \dots, W^Q(x, c_{Qj}) \leq W^Q(x^*, c_{Qj}) \end{array} \right\}.$$

$\mu_{\tilde{c}_p}(p = \overline{1, P})$ and $\mu_{\tilde{c}_q}(q = \overline{1, Q})$ are $(n \times 1)$ - ary α – level sets.

4 | Characterizing of α –Efficient Solutions for Problem (2)

To characterize the α – efficient solutions for *Problem (2)*, let us consider the following α – parametric MODNP problem:

$$\begin{aligned} \max Z^p(x, c_{pj}) &= \sum_{j=1}^n c_{pj} x_j, p = \overline{1, P}, \\ \min W^q(x, c_{qj}) &= \sum_{j=1}^n c_{qj} x_j, q = \overline{1, Q}. \end{aligned}$$

Subject to

$$x \in X, c_{pj} \in (\tilde{c}_{pj})_\alpha \text{ and } c_{qj} \in (\tilde{c}_{qj})_\alpha. \quad (6)$$

Where, $(\tilde{c}_{pj})$ and $(\tilde{c}_{qj})$ denote the α – level sets of the fuzzy variables $\tilde{c}_{pj}$ and $\tilde{c}_{qj}$. Based on the assumptions of the convexity, $\mu_{\tilde{c}_{pj}}, (\tilde{c}_{pj})_\alpha; \mu_{\tilde{c}_{qj}}, (\tilde{c}_{qj})_\alpha, (j = \overline{1, n}; p = \overline{1, P}; q = \overline{1, Q})$ are close intervals approximations of real numbers that are denoted by $[c_{pj}^-(\alpha), c_{pj}^+(\alpha)]$ and $[c_{qj}^-(\alpha), c_{qj}^+(\alpha)]$. Let χ_{pj}^α and χ_{qj}^α be the sets of $n \times 1$

matrices (c_{pj}) with $[c_{pj}^-(\alpha), c_{pj}^+(\alpha)]$ ($p = \overline{1, P}$) and (c_{qj}) with $[c_{qj}^-(\alpha), c_{qj}^+(\alpha)]$ ($q = \overline{1, Q}$). Problem (6) can be rewritten as

$$\begin{aligned} & \max Z^p(x, c_{pj}), p = \overline{1, P}, \\ & \min W^q(x, c_{qj}), q = \overline{1, Q}. \\ & \text{Subject to} \\ & x \in X, c_{pj} \in \chi_{pj}^\alpha \text{ and } c_{qj} \in \chi_{qj}^\alpha, p = \overline{1, P}; q = \overline{1, Q}. \end{aligned} \quad (7)$$

Theorem 1. $x^* \in X$ is said to be an α -efficient solution for Problem (2) if and only if $x^* \in X$ is an α -parametric efficient solution for Problem (6).

Proof: Necessity: Let $x^* \in X$ be an α -efficient solution for Problem (2) and $x^* \in X$ be not a α -n parametric efficient solution for Problem (6). Then there are $x^1 \in X, s_{pj} \in \chi_{pj}^\alpha$ and $t_{qj} \in \chi_{qj}^\alpha, p = \overline{1, P}; q = \overline{1, Q}; j = \overline{1, n}$. such that

$$Z^p(x, s_{pj}) \geq Z^p(x^*, s_{pj}) \text{ and } W^q(x, t_{qj}) \leq W^q(x^*, t_{qj}).$$

Since, $s_{pj} \in \chi_{pj}^\alpha$ and $t_{qj} \in \chi_{qj}^\alpha, p = \overline{1, P}; q = \overline{1, Q}; j = \overline{1, n}$. we have

$$\mu \left(\begin{aligned} & Z^1(x^1, \tilde{c}_{1j}) \geq Z^1(x^*, \tilde{c}_{1j}), \dots, Z^{p-1}(x^1, \tilde{c}_{p-1,j}) \geq Z^{p-1}(x^*, \tilde{c}_{p-1,j}), \\ & Z^p(x^1, \tilde{c}_{pj}) \geq Z^p(x^*, \tilde{c}_{pj}), Z^{p+1}(x^1, \tilde{c}_{p+1,j}) \geq Z^{p+1}(x^*, \tilde{c}_{p+1,j}), \dots, \\ & Z^P(x^1, \tilde{c}_{Pj}) \geq Z^P(x^*, \tilde{c}_{Pj}); W^1(x^1, \tilde{c}_{1j}) \leq W^1(x^*, \tilde{c}_{1j}), \dots, \\ & W^{q-1}(x^1, \tilde{c}_{q-1,j}) \leq W^{q-1}(x^*, \tilde{c}_{q-1,j}), W^q(x^1, \tilde{c}_{qj}) \leq W^q(x^*, \tilde{c}_{qj}), \\ & W^{q+1}(x^1, \tilde{c}_{q+1,j}) \leq W^{q+1}(x^*, \tilde{c}_{q+1,j}), \dots, W^Q(x^1, \tilde{c}_{Qj}) \leq W^Q(x^*, \tilde{c}_{Qj}) \end{aligned} \right) \geq \alpha.$$

Contradiction the assumption that $x^* \in X$ be an α -efficient solution for Problem (2).

Sufficiency: Let $x^* \in X$ be an α -parametric efficient solution for Problem (6) and $x^* \in X$ be not an α -efficient solution for Problem (2). Then, there are $x^1 \in X, p = \{1, \dots, P\}$ and $q = \{1, \dots, Q\}$ such that

$$\mu \left(\begin{aligned} & Z^1(x^2, \tilde{c}_{1j}) \geq Z^1(x^*, \tilde{c}_{1j}), \dots, Z^{p-1}(x^2, \tilde{c}_{p-1,j}) \geq Z^{p-1}(x^*, \tilde{c}_{p-1,j}), \\ & Z^p(x^2, \tilde{c}_{pj}) \geq Z^p(x^*, \tilde{c}_{pj}), Z^{p+1}(x^2, \tilde{c}_{p+1,j}) \geq Z^{p+1}(x^*, \tilde{c}_{p+1,j}), \dots, \\ & Z^P(x^2, \tilde{c}_{Pj}) \geq Z^P(x^*, \tilde{c}_{Pj}); W^1(x^2, \tilde{c}_{1j}) \leq W^1(x^*, \tilde{c}_{1j}), \dots, \\ & W^{q-1}(x^2, \tilde{c}_{q-1,j}) \leq W^{q-1}(x^*, \tilde{c}_{q-1,j}), W^q(x^2, \tilde{c}_{qj}) \leq W^q(x^*, \tilde{c}_{qj}), \\ & W^{q+1}(x^2, \tilde{c}_{q+1,j}) \leq W^{q+1}(x^*, \tilde{c}_{q+1,j}), \dots, W^Q(x^2, \tilde{c}_{Qj}) \leq W^Q(x^*, \tilde{c}_{Qj}) \end{aligned} \right) \geq \alpha, \quad (8)$$

i.e.,

$$\sup_{(c_1, \dots, c_P; c_1, \dots, c_Q) \in \hat{C} \times \hat{D}} \min \left(\begin{aligned} & \mu_{\tilde{c}_{1j}}(c_{1j}), \dots, \mu_{\tilde{c}_{p-1,j}}(c_{p-1,j}), \mu_{\tilde{c}_{pj}}(c_{pj}), \mu_{\tilde{c}_{p+1,j}}(c_{p+1,j}), \dots, \mu_{\tilde{c}_{Pj}}(c_{Pj}); \\ & \mu_{\tilde{c}_{1j}}(c_{1j}), \dots, \mu_{\tilde{c}_{q-1,j}}(c_{q-1,j}), \mu_{\tilde{c}_{qj}}(c_{qj}), \mu_{\tilde{c}_{q+1,j}}(c_{q+1,j}), \dots, \mu_{\tilde{c}_{Qj}}(c_{Qj}) \end{aligned} \right) \geq \alpha.$$

Where

$$\hat{C} = \left\{ \begin{aligned} & (c_1, \dots, c_P) \in \mathfrak{R}^{P \times Q(n \times 1)}; Z^1(x, c_{1j}) \geq Z^1(x^*, c_{1j}), \dots, \\ & Z^{p-1}(x, c_{p-1,j}) \geq Z^{p-1}(x^*, c_{p-1,j}), Z^p(x, c_{pj}) \geq Z^p(x^*, c_{pj}), \\ & Z^{p+1}(x, c_{p+1,j}) \geq Z^{p+1}(x^*, c_{p+1,j}), \dots, Z^P(x, c_{Pj}) \geq Z^P(x^*, c_{Pj}). \end{aligned} \right\}.$$

And

$$\widehat{D} = \left\{ \begin{array}{l} (c_1, \dots, c_Q) \in \mathfrak{R}^{Q(n \times 1)}: W^1(x^2, c_{1j}) \leq W^1(x^*, c_{1j}), \dots, \\ W^{q-1}(x^2, c_{q-1,j}) \leq W^{q-1}(x^*, c_{q-1,j}), W^q(x^2, c_{qj}) \leq W^q(x^*, c_{qj}), \\ W^{q+1}(x^2, c_{q+1,j}) \leq W^{q+1}(x^*, c_{q+1,j}), \dots, W^Q(x^2, c_{Qj}) \leq W^Q(x^*, c_{Qj}). \end{array} \right\}.$$

In the case for existing the supremum, there is $(u_{1j}, \dots, u_{pj}) \in \widehat{C}$ and $(v_{1j}, \dots, v_{Qj}) \in \widehat{D}$ with

$\min\{\mu_{\tilde{c}_{1j}}(u_{1j}), \dots, \mu_{\tilde{c}_{pj}}(u_{pj})\} < \alpha$, and $\min\{\mu_{\tilde{c}_{1j}}(v_{1j}), \dots, \mu_{\tilde{c}_{Qj}}(v_{Qj})\} < \alpha$, then

$$\sup_{(c_1, \dots, c_P; c_1, \dots, c_Q) \in \widehat{C} \times \widehat{D}} \min \left(\begin{array}{l} \mu_{\tilde{c}_{1j}}(c_{1j}), \dots, \mu_{\tilde{c}_{p-1,j}}(c_{p-1,j}), \mu_{\tilde{c}_{pj}}(c_{pj}), \mu_{\tilde{c}_{p+1,j}}(c_{p+1,j}), \dots, \mu_{\tilde{c}_{pj}}(c_{pj}); \\ \mu_{\tilde{c}_{1j}}(c_{1j}), \dots, \mu_{\tilde{c}_{q-1,j}}(c_{q-1,j}), \mu_{\tilde{c}_{qj}}(c_{qj}), \mu_{\tilde{c}_{q+1,j}}(c_{q+1,j}), \dots, \mu_{\tilde{c}_{Qj}}(c_{Qj}) \end{array} \right) < \alpha.$$

This is a contradiction (4). Then there are $(u_{1j}, \dots, u_{pj}) \in \widehat{C}$ and $(v_{1j}, \dots, v_{Qj}) \in \widehat{D}$ with

$\min\{\mu_{\tilde{c}_{1j}}(u_{1j}), \dots, \mu_{\tilde{c}_{pj}}(u_{pj})\} \geq \alpha$, and $\min\{\mu_{\tilde{c}_{1j}}(v_{1j}), \dots, \mu_{\tilde{c}_{Qj}}(v_{Qj})\} \geq \alpha$, i. e.,

$$u_{pj} \in \chi_{pj}^\alpha (p = \overline{1, P}) \text{ and } v_{qj} \in \chi_{qj}^\alpha (q = \overline{1, Q}). \quad (9)$$

From Eqs. (4) and (10), we conclude that $x^* \in X$ is an α -parametric efficient solution for Problem (6).

5 | Min-max GP Approach

α -Parametric MODNP Problem (6) can be demonstrated using a min-max GP approach as $\min d$

Subject to (10)

$$Z^p(x, c_{pj}) + l_p - f_p = Z^{p*}, l_p \leq d,$$

$$\gamma_p \frac{l_p}{y_p} \leq d, y_p = Z^{p*} - Z^{p-}.$$

$$W^q(x, c_{qj}) + l_q - f_q = W^{q*},$$

$$\delta_q \frac{d_q}{y_q} \leq d, W^{q-} - W^{q*}.$$

$$Vx \leq B,$$

$$c_{pj} \in [c_{pj}^-(\alpha), c_{pj}^+(\alpha)], c_{qj} \in [c_{qj}^-(\alpha), c_{qj}^+(\alpha)], p = \overline{1, P}; q = \overline{1, Q}; j = \overline{1, n}, 0 \leq d \leq 1.$$

Where, $Z^{p*} = \max Z^p$, and $W^{q*} = \min W^q$, are the positive ideal solutions, respectively. Also, $Z^{p*} = \min Z^p$, and $W^{q*} = \max W^q$ are the negative ideal solutions; respectively, γ_p and δ_q are positive weights, $y_p = Z^{p*} - Z^{p-}$, and $y_q = W^{q-} - W^{q*}$ are the normalization of the positive and negative ideal solutions, respectively.

Now, let us determine the stability set of the first kind $S(\hat{x}, \hat{c})$ corresponding to the α -optimal solution by applying the following condition:

$$\theta_{pj}(\hat{c}_{pj} - c_{pj}^+) = 0, p = \overline{1, P}; j = \overline{1, n},$$

$$\theta_{pj}(c_{pj}^- - \hat{c}_{pj}) = 0, p = \overline{1, P}; j = \overline{1, n},$$

$$\theta_{qj}(\hat{c}_{qj} - c_{qj}^+) = 0, q = \overline{1, Q}; j = \overline{1, n},$$

$$\theta_{qj}(c_{qj}^- - \hat{c}_{qj}) = 0, q = \overline{1, Q}; j = \overline{1, n}.$$

Where

$$[c_{pj}^-(\alpha), c_{pj}^+(\alpha)] = L_\alpha(\tilde{c}_{pj}), p = \overline{1, P}, j = \overline{1, n}, \text{ and } [c_{qj}^-(\alpha), c_{qj}^+(\alpha)], q = \overline{1, Q}.$$

Consider the following cases:

$$I. \theta_{pj} > 0, p \in J_1 \subset \{1, \dots, P\}; \theta_{pj} = 0, p \notin J_1,$$

$$\theta_{qj} > 0, q \in J_1 \subset \{1, \dots, Q\}; \theta_{qj} = 0, q \notin J_1,$$

$$\theta_{pj} > 0, i \in J_2 \subset \{1, \dots, P\}, \theta_{pj} = 0, i \notin J_2,$$

$$\theta_{qj} > 0, i \in J_2 \subset \{1, \dots, Q\}, \theta_{qj} = 0, i \notin J_2.$$

Let M be the set of all proper subsets of $\{1, \dots, P\}$ and $\{1, \dots, Q\}$. Then

$$S_{J_1, J_2}(\hat{x}, \hat{c}) = \left\{ \begin{array}{l} \{(c_{pj}^-, c_{pj}^+), (c_{qj}^-, c_{qj}^+)\} \in \mathfrak{R}^{2p \times 2q}: c_{pj}^+ = \hat{c}_{pj}, p \in J_1, c_{pj}^+ \geq \hat{c}_{pj}, p \notin J_1, \\ c_{qj}^+ = \hat{c}_{qj}, q \in J_1, c_{qj}^+ \geq \hat{c}_{qj}, q \notin J_1; c_{pj}^- = \hat{c}_{pj}, p \in J_2, c_{pj}^- \leq \hat{c}_{pj}, p \notin J_2, \\ c_{qj}^- = \hat{c}_{qj}, q \in J_2, c_{qj}^- \leq \hat{c}_{qj}, q \notin J_2. \end{array} \right\}.$$

Hence

$$S_1(\hat{x}, \hat{c}) = \bigcup_{J_1, J_2 \in M} S_{J_1, J_2}(\hat{x}, \hat{c}).$$

II. $\theta_{pj}, \theta_{pj}; \theta_{qj}, \theta_{qj} = 0$, then

$$S_2(\hat{x}, \hat{c}) = \left\{ \begin{array}{l} \{(c_{pj}^-, c_{pj}^+), (c_{qj}^-, c_{qj}^+)\} \in \mathfrak{R}^{2p \times 2q}: c_{pj}^+ \geq \hat{c}_{pj}, p = \overline{1, P}, \\ c_{qj}^+ \geq \hat{c}_{qj}, q = \overline{1, Q}; c_{pj}^- \leq \hat{c}_{pj}, p = \overline{1, P}, c_{qj}^- \leq \hat{c}_{qj}, q = \overline{1, Q} \end{array} \right\}.$$

III. $\theta_{pj}, \theta_{pj}; \theta_{qj}, \theta_{qj} > 0$, then

$$S_3(\hat{x}, \hat{c}) = \left\{ \begin{array}{l} \{(c_{pj}^-, c_{pj}^+), (c_{qj}^-, c_{qj}^+)\} \in \mathfrak{R}^{2p \times 2q}: c_{pj}^+ = \hat{c}_{pj}, p = \overline{1, P}, \\ c_{qj}^+ = \hat{c}_{qj}, q = \overline{1, Q}; c_{pj}^- = \hat{c}_{pj}, p = \overline{1, P}, c_{qj}^- = \hat{c}_{qj}, q = \overline{1, Q} \end{array} \right\}.$$

Hence

$$S(\hat{x}, \hat{c}) = \bigcup_{k=1}^K S_k(\hat{x}, \hat{c}).$$

6 | Numerical Example

Consider the following problem

- I. $\max \tilde{Z}^1 = \tilde{c}_{11}x_1 + \tilde{c}_{12}x_2 + \tilde{c}_{13}x_3$, (Profits).
- II. $\max \tilde{Z}^2 = \tilde{c}_{21}x_1 + \tilde{c}_{22}x_2 + \tilde{c}_{23}x_3$, (Quality).
- III. $\max \tilde{Z}^3 = \tilde{c}_{31}x_1 + \tilde{c}_{32}x_2 + \tilde{c}_{33}x_3$, (Workers Satisfaction).

Subject to

$$12x_1 + 17x_2 \leq 1400, \text{ (Milling Machine).}$$

$$3x_1 + 9x_2 + 8x_3 \leq 1000, \text{ (Lathe).}$$

$$10x_1 + 13x_2 + 15x_3 \leq 1750, \text{ (Grinder).}$$

$$6x_1 + 0x_2 + 16x_3 \leq 1325, \text{ (Jig Saw).}$$

$$0x_1 + 12x_2 + 7x_3 \leq 900, \text{ (Drill Press).}$$

$$x_1, x_2, x_3 \geq 0.$$

With the price of resources $p_1 = \$ 0.75, p_2 = \$ 0.6, p_3 = \$ 0.35, p_4 = \$ 0.50, p_5 = \$ 1.15$ and $p_6 = \$ 0.65$, and the budget level $B = \$ 4658.75$,

$$\tilde{c}_{11} = (20, 30, 50, 60, 80), \tilde{c}_{12} = (70, 85, 100, 110, 130), \tilde{c}_{13} = (13.5, 15.5, 17.5, 20.5, 22.5),$$

$$\tilde{c}_{21} = (85, 90, 92, 97, 105), \tilde{c}_{22} = (60, 70, 75, 85, 100), \tilde{c}_{23} = (30, 45, 50, 60, 75),$$

$$\tilde{c}_{31} = (10, 18, 25, 30, 45), \tilde{c}_{32} = (80, 90, 100, 110, 115), \tilde{c}_{33} = (55, 75, 75, 80, 95).$$

Problem (11) can be formulated according to *Problem (2)* as

(11)

$$\begin{aligned}
 \max \tilde{Z}^1 &= \tilde{c}_{11}x_1 + \tilde{c}_{12}x_2 + \tilde{c}_{13}x_3, \text{ (Profits).} \\
 \max \tilde{Z}^2 &= \tilde{c}_{21}x_1 + \tilde{c}_{22}x_2 + \tilde{c}_{23}x_3, \text{ (Quality).} \\
 \max \tilde{Z}^3 &= \tilde{c}_{31}x_1 + \tilde{c}_{32}x_2 + \tilde{c}_{33}x_3, \text{ (Workers Satisfaction).} \\
 \text{Subject to} \\
 23.475x_1 + 42.675x_2 + 28.7x_3 &= 4558.75, \\
 x_1, x_2, x_3 &\geq 0.
 \end{aligned} \tag{12}$$

The positive ideal solutions to *Problem (11)* are

$$\begin{aligned}
 \max Z^1 &= 50x_1 + 100x_2 + 17.5x_3, \\
 \max Z^2 &= 92x_1 + 75x_2 + 50x_3, \\
 \max Z^3 &= 25x_1 + 100x_2 + 75x_3. \\
 \text{Subject to} \\
 23.475x_1 + 42.675x_2 + 28.7x_3 &= 4558.75, \\
 x_1, x_2, x_3 &\geq 0.
 \end{aligned} \tag{13}$$

Table 1. Positive ideal solutions to problem (13).

Decision Variables	Z^1	Z^2	Z^3
x_1	0	194.1960	0
x_2	106.8248	0	0
x_3	0	0	158.8415
	$\max Z^{1*} = 10682.48$	$\max Z^{2*} = 17866.03$	$\max Z^{3*} = 11913.11$

The negative ideal solutions to *Problem (11)* are.

$$\begin{aligned}
 \min Z^1 &= 20x_1 + 130x_2 + 13.5x_3. \\
 \min Z^2 &= 105x_1 + 60x_2 + 75x_3. \\
 \min Z^3 &= 10x_1 + 115x_2 + 55x_3. \\
 \text{Subject to} \\
 23.475x_1 + 42.675x_2 + 28.7x_3 &= 4558.75, \\
 x_1, x_2, x_3 &\geq 0.
 \end{aligned} \tag{14}$$

Table 2. Negative ideal solutions to problem (14).

Decision variables	Z^1	Z^2	Z^3
x_1	0	0	194.1960
x_2	0	106.8248	0
x_3	158.8415	0	0
	$\max Z^{1*} = 2144.360$	$\max Z^{2*} = 6409.490$	$\max Z^{3*} = 1941.960$

Based on the positive and negative ideal solutions, *Problem (11)*, referring to *Problem (10)*, becomes min d subject to

$$\begin{aligned}
 c_{11}x_1 + c_{12}x_2 + c_{13}x_3 + l_1 - f_1 &= 10682.48, \\
 c_{21}x_1 + c_{22}x_2 + c_{23}x_3 + l_2 - f_2 &= 17866.03, \\
 c_{31}x_1 + c_{32}x_2 + c_{33}x_3 + l_3 - f_3 &= 11913.11, \\
 \frac{l_1}{10682.48 - 2144.360} &\leq d, \\
 \frac{l_2}{17866.03 - 6409.490} &\leq d, \\
 \frac{l_3}{11913.11 - 1941.960} &\leq d, \\
 23.475x_1 + 42.675x_2 + 28.7x_3 &= 4558.75, \\
 x_1, x_2, x_3 &\geq 0, 0 \leq d \leq 1, \\
 c_{11} &= [30, 60], c_{12} = [85, 110], c_{13} = [15.5, 20.5], \\
 c_{21} &= [90, 97], c_{22} = [70, 85], c_{23} = [45, 65], \\
 c_{31} &= [18, 30], c_{32} = [90, 110], c_{33} = [75, 80].
 \end{aligned} \tag{15}$$

Table 3. An optimal satisfactory solution to problem (15).

Decision Variables	Values
$\hat{x}_1$	89
$\hat{c}_{11}$	60
$\hat{c}_{12}$	110
$\hat{c}_{13}$	20.5
$\hat{x}_2$	17.311
$\hat{c}_{21}$	97
$\hat{c}_{22}$	85
$\hat{c}_{23}$	80
$\hat{x}_3$	59.76241
$\hat{c}_{31}$	30
$\hat{c}_{32}$	110
$\hat{c}_{33}$	65
l_1	2173.183
l_1	2916
l_1	2537.928
f_1	0
f_1	0
f_1	0
$\max \widehat{Z}^1$	= 8529.34
$\max \widehat{Z}^2$	= 14885.43
$\max \widehat{Z}^3$	= 8458.77
d	= 0.2545271

It is clear that the results obtained by the proposed method are less than those obtained by Umarusman [13].

Hence, $S(\hat{x}, \hat{c})$ is determined by applying the following conditions:

$$\vartheta_{1j}(\hat{c}_{1j} - c_{1j}^+) = 0, j = 1, 2, 3,$$

$$\vartheta_{2j}(\hat{c}_{1j} - c_{1j}^+) = 0, j = 1, 2, 3,$$

$$\vartheta_{3j}(\hat{c}_{1j} - c_{1j}^+) = 0, j = 1, 2, 3, \vartheta_{1j}, \vartheta_{2j}, \vartheta_{3j} \geq 0.$$

We have $J_1 \subseteq \{1, 2, 3\}$. For $J_1 = \emptyset, \vartheta_{1j} = \vartheta_{2j} = \vartheta_{3j} = 0, j = 1, 2, 3$. Then

$$S_{J_1}(\hat{x}, \hat{c}) = \left\{ c \in \mathfrak{R}^9 : c_{11} \geq 60, c_{12} \geq 110, c_{13} \geq 20.5, c_{21} \geq 97, \right. \\ \left. c_{22} \geq 85, c_{23} \geq 80, c_{31} \geq 30, c_{32} \geq 110, c_{33} \geq 65 \right\}.$$

For $J_2 = \{1\}, \vartheta_{1j} > 0, \vartheta_{2j} = 0, \vartheta_{3j} = 0$. Then

$$S_{J_2}(\hat{x}, \hat{c}) = \left\{ c \in \mathfrak{R}^9 : c_{11} = 60, c_{12} = 110, c_{13} = 20.5, c_{21} \geq 97, \right. \\ \left. c_{22} \geq 85, c_{23} \geq 80, c_{31} = 30, c_{32} = 110, c_{33} = 65 \right\}.$$

For $J_3 = \{2\}, \vartheta_{1j} = 0, \vartheta_{2j} > 0, \vartheta_{3j} = 0$. Then

$$S_{J_3}(\hat{x}, \hat{c}) = \left\{ c \in \mathfrak{R}^9 : c_{11} \geq 60, c_{12} \geq 110, c_{13} \geq 20.5, c_{21} = 97, \right. \\ \left. c_{22} = 85, c_{23} = 80, c_{31} \geq 30, c_{32} \geq 110, c_{33} \geq 65 \right\}.$$

For $J_4 = \{3\}, \vartheta_{1j} = 0, \vartheta_{2j} = 0, \vartheta_{3j} > 0$. Then

$$S_{J_4}(\hat{x}, \hat{c}) = \left\{ c \in \mathfrak{R}^9 : c_{11} \geq 60, c_{12} \geq 110, c_{13} \geq 20.5, c_{21} \geq 97, \right. \\ \left. c_{22} \geq 85, c_{23} \geq 80, c_{31} = 30, c_{32} = 110, c_{33} = 65 \right\}.$$

For $J_5 = \{1, 2\}, \vartheta_{1j} > 0, \vartheta_{2j} > 0, \vartheta_{3j} = 0$. Then

$$S_{J_5}(\hat{x}, \hat{c}) = \left\{ c \in \mathfrak{R}^9 : c_{11} = 60, c_{12} = 110, c_{13} = 20.5, c_{21} = 97, \right. \\ \left. c_{22} = 85, c_{23} = 80, c_{31} \geq 30, c_{32} \geq 110, c_{33} \geq 65 \right\},$$

For $J_6 = \{1, 3\}, \vartheta_{1j} > 0, \vartheta_{2j} = 0, \vartheta_{3j} > 0$. Then

$$S_{J_6}(\hat{x}, \hat{c}) = \left\{ c \in \mathfrak{R}^9 : c_{11} \geq 60, c_{12} \geq 110, c_{13} \geq 20.5, c_{21} = 97, \right. \\ \left. c_{22} = 85, c_{23} = 80, c_{31} \geq 30, c_{32} \geq 110, c_{33} \geq 65 \right\}.$$

For $J_7 = \{2, 3\}, \vartheta_{1j} = 0, \vartheta_{2j} > 0, \vartheta_{3j} > 0$. Then

$$S_{J_7}(\hat{x}, \hat{c}) = \left\{ c \in \mathfrak{R}^9 : c_{11} \geq 60, c_{12} \geq 110, c_{13} \geq 20.5, c_{21} = 97, \right. \\ \left. c_{22} = 85, c_{23} = 80, c_{31} = 30, c_{32} = 110, c_{33} = 65 \right\}.$$

For $J_8 = \{1, 2, 3\}, \vartheta_{1j} > 0, \vartheta_{2j} > 0, \vartheta_{3j} > 0$. Then

$$S_{j_8}(\hat{x}, \hat{c}) = \left\{ c \in \mathbb{R}^9 : c_{11} = 60, c_{12} = 110, c_{13} = 20.5, c_{21} = 97, \right. \\ \left. c_{22} = 85, c_{23} = 80, c_{31} = 30, c_{32} = 110, c_{33} = 65 \right\}$$

Hence

$$S(\hat{x}, \hat{c}) = \bigcup_{k=1}^8 S_k(\hat{x}, \hat{c}).$$

7 | Conclusions

The De novo hypothesis provides meta0 optimum solutions at optimal levels for single and multi-objective programming. This paper studies DNP in an uncertain environment. Close interval approximation of the PQF number is applied to solve the MODNP problem. A necessary and sufficient condition for the solution from the efficiency standpoint has been established.

A Min-max goal programming approach with positive and negative ideals has been proposed for optimal compromise system design. The stability set of the first kind corresponding to the optimal system design has been defined and determined. The advantages of this approach are that it can be applied to any environment and enables the decision-maker to investigate real-world problems.

Author Contributions

Hamiden Abd El-Wahed Khalifa conceptualized and designed the study, developed the theoretical framework, and led the analysis and interpretation of the data. Seyed Ahmad Edalatpanah contributed to the development of the mathematical models and provided insights into the fuzzy optimization techniques applied in the study. Sultan S. Alodhaibi assisted with the implementation of the numerical experiments, data collection, and the computational analysis. All authors contributed to the writing, reviewing, and final approval of the manuscript.

Data Availability

The datasets generated and/or analyzed during the current study are available from the corresponding author on reasonable request.

Conflicts of Interest

The authors declare no conflict of interest related to the publication of this paper.

References

- [1] Hwang, C. L., Paidy, S. R., Masud, A. S. M., & Yoon, K. (2012). *Multiple objective decision making — methods and applications: a state-of-the-art survey*. Springer Berlin Heidelberg. <https://books.google.com/books?id=M0noCAAAQBAJ>
- [2] Wang, S., & Chiang, H. D. (2019). Constrained multiobjective nonlinear optimization: a user preference enabling method. *IEEE transactions on cybernetics*, 49(7), 2779–2791. DOI:10.1109/TCYB.2018.2833281
- [3] Kundu, T., & Islam, S. (2019). An interactive weighted fuzzy goal programming technique to solve multi-objective reliability optimization problem. *Journal of industrial engineering international*, 15(1), 95–104. <https://doi.org/10.1007/s40092-019-0321-y>
- [4] Waliv, R. H., Mishra, U., Garg, H., & Umap, H. P. (2020). A nonlinear programming approach to solve the stochastic multi-objective inventory model using the uncertain information. *Arabian journal for science and engineering*, 45(8), 6963–6973. <https://doi.org/10.1007/s13369-020-04618-z>

- [5] Ahmad, F. (2021). Robust neutrosophic programming approach for solving intuitionistic fuzzy multiobjective optimization problems. *Complex & intelligent systems*, 7(4), 1935–1954. <https://doi.org/10.1007/s40747-021-00299-9>
- [6] Liu, X., Liu, Y., He, X., Xiao, M., & Jiang, T. (2021). Multi-objective nonlinear programming model for reducing octane number loss in gasoline refining process based on data mining technology. *Processes*, 9(4), 1–19. <https://www.mdpi.com/2227-9717/9/4/721>
- [7] Zeleny, M. (2012). *Multiple criteria decision making kyoto 1975*. Springer Berlin Heidelberg. <https://books.google.com/books?id=kUrmCAAQBAJ>
- [8] Zeleny, M. (1986). Optimal system design with multiple criteria: de novo programming approach. *Engineering costs and production economics*, 10(1), 89–94. <https://www.sciencedirect.com/science/article/pii/0167188X86900297>
- [9] Zelený, M. (1990). Optimizing given systems vs. designing optimal systems: the de novo programming approach. *International journal of general systems*, 17(4), 295–307. <https://doi.org/10.1080/03081079008935113>
- [10] Shi, Y. (1999). Optimal system design with multiple decision makers and possible debt: a multicriteria de novo programming approach. *Operations research*, 47(5), 723–729. <https://doi.org/10.1287/opre.47.5.723>
- [11] Zhang, Y. M., Huang, G. H., & Zhang, X. D. (2009). Inexact de novo programming for water resources systems planning. *European journal of operational research*, 199(2), 531–541. <https://www.sciencedirect.com/science/article/pii/S0377221708010060>
- [12] Chen, J. K. C., & Tzeng, G. H. (2009). Perspective strategic alliances and resource allocation in supply chain systems through the de novo programming approach. *International journal of sustainable strategic management*, 1(3), 320–339. <https://www.inderscienceonline.com/doi/abs/10.1504/IJSSM.2009.026285>
- [13] Umarusman, N. (2013). Min-max goal programming approach for solving multi-objective de novo programming problems. *International journal of operations research*, 10(2), 92–99. <https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=4dd384b06495c580c325ef9d262fb86266d655cb>
- [14] Zadeh, L. A. (1965). Fuzzy sets. *Information and control*, 8(3), 338–353. <https://www.sciencedirect.com/science/article/pii/S00199586590241X>
- [15] Dubois, D. J. (1980). *Fuzzy sets and systems: theory and applications*. Elsevier Science. <https://books.google.com/books?id=JmjfHUUtMkMC>
- [16] Bellman, R. E., & Zadeh, L. A. (1970). Decision-making in a fuzzy environment. *Management science*, 17(4), 1–141. <https://doi.org/10.1287/mnsc.17.4.B141>
- [17] Zimmermann, H. J. (1978). Fuzzy programming and linear programming with several objective functions. *Fuzzy sets and systems*, 1(1), 45–55. <https://www.sciencedirect.com/science/article/pii/0165011478900313>
- [18] Chen, Y. W., & Hsieh, H. E. (2006). Fuzzy multi-stage de-novo programming problem. *Applied mathematics and computation*, 181(2), 1139–1147. <https://www.sciencedirect.com/science/article/pii/S0096300306001986>
- [19] J-J Huang, G. H. T., & Ong, C. S. (2006). Choosing best alliance partners and allocating optimal alliance resources using the fuzzy multi-objective dummy programming model. *Journal of the operational research society*, 57(10), 1216–1223. <https://doi.org/10.1057/palgrave.jors.2602108>
- [20] Charnes, A., & Cooper, W. W. (1957). Management models and industrial applications of linear programming. *Management science*, 4(1), 38–91. <https://doi.org/10.1287/mnsc.4.1.38>
- [21] Lee, S. M. (1972). *Goal programming for decision analysis*. Auerbach Publishers. <https://cir.nii.ac.jp/crid/1130000794581506944>
- [22] Ijiri, Y. (1965). *Management goals and accounting for control*. North Holland Publishing Company. <https://books.google.com/books?id=KCKUAQAAMAAJ>
- [23] Ignizio, J. P. (1982). *Linear programming in single- multiple-objective systems*. Prentice-Hall. <https://cir.nii.ac.jp/crid/1130282271612104960>
- [24] Jones, D., Tamiz, M., & others. (2010). *Practical goal programming* (Vol. 141). Springer. <https://link.springer.com/content/pdf/10.1007/978-1-4419-5771-9.pdf>
- [25] Osman, M. S. A. (1977). Qualitative analysis of basic notions in parametric convex programming. I. Parameters in the constraints. *Aplikace matematiky*, 22(5), 318–332. <https://dml.cz/dmlcz/103710>

- [26] Osman, M. S., & El-Banna, A. Z. H. (1993). Stability of multiobjective nonlinear programming problems with fuzzy parameters. *Mathematics and computers in simulation*, 35(4), 321–326.
<https://www.sciencedirect.com/science/article/pii/037847549390062Y>
- [27] Jain, S. (2010). Close interval approximation of piecewise quadratic fuzzy numbers for fuzzy fractional program. *Iranian journal of operations research*, 2(1), 77–88.
https://www.sid.ir/EN/VEWSSID/J_pdf/115720100106.pdf
- [28] Chankong, V., & Haimes, Y. Y. (2008). *Multiobjective decision making: theory and methodology*. Dover Publications. <https://books.google.com/books?id=o371DAAAQBAJ>